

INTEGRATING BIODIVERSITY AND ENVIRONMENTAL OBSERVATIONS

IN SUPPORT OF NATIONAL MARINE SANCTUARY AND LARGE MARINE ECOSYSTEM ASSESSMENTS

By Henry A. Ruhl, Jennifer A. Brown, Alexandra R. Harper, Elliott L. Hazen, Lynn deWitt, Patrick Daniel, Andrew DeVogelaere, Raphael M. Kudela, John P. Ryan, Alexis D. Fischer, Frank E. Muller-Karger, and Francisco P. Chavez



Young-of-the-year vermilion rockfish (*Sebastes miniatus*) at Dali's Wall in Carmel Bay, Monterey Bay National Marine Sanctuary. Photo credit: Steve Lonhart/NOAA MBNMS

ABSTRACT. Species and habitats are the subjects of legislation that mandates reporting of information on ecosystem conditions. Improvements in sensors, sampling platforms, information systems, and collaborations among experts and information users now enables more effective and up-to-date information to meet regional and national needs. Specifically, advances in environmental DNA (eDNA)-based assessments of biodiversity, community science data, various underwater imaging devices, and environmental, behavioral, and physiology observations from animal telemetry provide new opportunities to address multiple requirements for reporting status and trends, including insights into life in the deep ocean. Passive and active acoustic sensors help monitor marine life, boat traffic, and noise pollution. Satellites provide repeated, frequent, and long-term records of many relevant variables from global to local scales and, when combined with numerical computer simulations, allow planning for future scenarios. Metadata standards facilitate the transfer of data from machine to machine, thus streamlining assessments and forecasting and providing knowledge directly to the public. The Marine Biodiversity Observation Network (MBON) facilitates this exchange of information on life in the sea. The collaborative efforts of the Central and Northern California Ocean Observing System (CeNCOOS) of the US Integrated Ocean Observing System and its partners provide an example of a regional MBON process for information delivery. This includes linking policy and management needs, prioritizing observing data from various platforms and methods, streamlining data handling practices, and delivering information for management such as for the Monterey Bay National Marine Sanctuary and the California Current Large Marine Ecosystem, with iterative process adaptation.

INTRODUCTION

Managers of coastal and marine resources need the most current information on changing marine life and environmental conditions. They are responsible for protecting the public and natural resources and are required to meet national, state, and local regulatory and reporting requirements. Biological and biodiversity information is required in US legislation, such as the Magnuson-Stevens Act, the Marine Mammal Protection Act, the National Marine Sanctuaries Act, the Endangered Species Act, the Coordinated Ocean Observations and Research Act of 2020, and in many other statutes from international to state and local levels (Hildreth, 2008; Bax et al., 2019). Ecosystem observations are the cornerstones of marine conservation, management, science research, and education programs. Healthy marine ecosystems form the basis of the livelihood and well-being of numerous communities and industries globally (Estes et al., 2021). Yet biological and ecosystem level observations are often difficult to access, are costly, and are presented in multiple different formats.

A new paradigm, in which data from multiple groups for multiple purposes are shared and are more rapidly available, is emerging with efforts such as the US Marine Biodiversity Observation Network (MBON). Improved delivery of ecosystem assessment information is facilitating Ecosystem Status Reports (ESRs) of the Integrated Ecosystem Assessment (IEA) Program (Harvey et al., 2020; Monaco et al., 2021) and Condition Reports of the National Oceanic and Atmospheric Administration (NOAA) Office of National Marine Sanctuaries. Here, we focus on examples of progress for the California Current Large Marine Ecosystem (CCLME) and Monterey Bay National Marine Sanctuary (MBNMS). Several MBON initiatives integrate existing efforts, advance new modes of observing, and create a scalable framework for addressing regulatory directives. To realize the vision of integrated observing for biology and biodiversity, the MBON program seeks to be integrative and to build new partnerships, many of which are developed around in-kind contributions and the mutual support required

to develop common best practices for solving problems through standardized observation methods and applications.

Best practices adopted by MBON contributors are allowing the integration of several forms of data into a common infrastructure. Interoperable data facilitates the development of assessments for National Marine Sanctuaries, California marine protected areas (MPAs), oil and gas development proposals, and proposed wind energy farms. Challenges to delivering this information efficiently include the time required to quality control observations and bringing data into comparable formats. In many cases, format requirements for analytics are not followed by observers, delaying analyses and delivery of information to resource managers.

Within the Central and Northern California region, MBON includes regional partners such as the Central and Northern California Ocean Observing System (CeNCOOS), national partners including the Integrated Ocean Observing System (IOOS), and global partners such as the Group on Earth Observations Biodiversity Observation Network (GEO BON) and the Global Ocean Observing System (GOOS). Individuals from around the world are sharing their data and experiences to reuse and add value to biology and biodiversity information. Here, we describe existing and maturing “building blocks” for improved coordination of the wealth of data collected by these efforts.

REGIONAL INTRODUCTION AND MBON PROJECT BACKGROUND

The California Current Large Marine Ecosystem ranges from Baja California to the northern tip of Vancouver Island and encompasses diverse and dynamic oceanic and coastal habitats. It is strongly influenced by seasonal upwelling that supports high primary productivity and fish abundance from the coast to areas far offshore over the deep ocean (e.g., Chavez et al., 2003, 2017; Ruhl and Smith, 2004; Woodson et al., 2012). These phenomena are strongly modulated by interannual variations in basin circulation, such as the

El Niño-Southern Oscillation (ENSO), as well as by longer-term variability such as the Pacific Decadal Oscillation (PDO) and the North Pacific Gyre Oscillation (NPGO) (e.g., Mantua et al., 1997; Schwing et al., 2002; Di Lorenzo et al., 2008). In 2014–2016, a large area of the Northeast Pacific became known as the “Warm Blob” (e.g., Bond et al., 2015; García-Reyes and Sydeman, 2017). Such large-scale phenomena can exert a strong influence on marine ecosystem changes (Chavez et al. 2017; Ryan et al., 2017; Barth et al., 2018). On top of such large-scale changes, there is considerable along-shore variability in ecosystem conditions, such as spatial differences in upwelling and CO₂ fluxes associated with coastal promontories (Chavez and Messie, 2009; Fiechter et al., 2014).

The Monterey Bay region, located in the south-central portion of the CCLME, is renowned for its marine biodiversity. However, considerable work is needed to synchronize, coordinate, and make biological observing efforts interoperable across the large number of marine science and conservation organizations in the region that represent academic, federal, state, private, and community sectors. In 2014, the Bureau of Ocean Energy Management (BOEM), the National Aeronautics and Space Administration (NASA), and others funded a series of MBON projects through the National Oceanographic Partnership Program (NOPP), including an early project focusing on the central and northern California region and the Florida Keys titled National Marine Sanctuaries as Sentinel Sites for a Demonstration Marine Biodiversity Observation Network (Sanctuaries MBON). Completed in 2020, this project was a collaborative effort between the MBNMS, members of the CCLME IEA team at the NOAA National Marine Fisheries Service Southwest Fisheries Science Center (SWFSC), CeNCOOS, and other regional partners. This Sanctuaries MBON was designed to define essential variables for biological monitoring (Muller-Karger et al., 2018) and

to develop eDNA method and application pipelines (Chavez et al., 2021, in this issue), diversity and species richness indices (Santora et al., 2017), assessments of changing biogeographical seascapes (Kavanaugh et al., 2021, in this issue), and data management and information tools to underpin delivery of each of these (e.g., Benson et al., 2021, in this issue), and to provide information for Monterey Bay and Florida Keys National Marine Sanctuaries. These efforts converged on the modern and flexible Darwin Core data standard for biological taxonomic data for operational and research applications. This led to more biological data being made available for public access through automated servers (e.g., a NOAA SWFSC ERDDAP data server; Simons, 2019).

A new MBON project in central California was initiated in 2019 to focus on the quantification of relationships among climate, ocean physics, biogeochemistry, and ecosystem dynamics affecting everything from microbes to top predators. Key aims included improvement of forecasts for forage fish distributions and evaluations of the physical and biological characteristics of the satellite-based seascapes biogeographic classification system (Kavanaugh et al., 2021, in this issue). The CCLME IEA and MBNMS Condition Reports are key delivery targets for the information developed by MBON efforts. The “Central California MBON” complements other regional MBON projects that address similar requirements in southern California, the Pacific Northwest, the Arctic, the Gulf of Maine, and South Florida. These US MBON initiatives, in turn, join projects globally contributing to GEO BON and GOOS.

UNDERSTANDING NEEDS AND CO-DESIGNING SOLUTIONS

The US IOOS program, implemented in central/northern California through the CeNCOOS Regional Association, operates and maintains observing systems, manages data, and adds value through data and information product develop-

ment, and engages stakeholders regularly (CeNCOOS, 2020). This work has included regional development of technology, methods, and research to understand processes and theory. A substantial focus is working with current and future observing system contributors and the public. Serving diverse stakeholder needs requires communication, consultation, and collaborative approaches—including public-private partnerships. CeNCOOS works with managers at NOAA’s National Marine Sanctuaries and Pacific Fishery Management Council, the California Ocean Protection Council (OPC), and the California Departments of Fish and Wildlife (CDFW), Water (CDW), and Public Health (CDPH) to make the data accessible in formats that meet the needs of users with a variety of technical abilities.

ONMS Condition Reports and Information Needs for Management

NOAA’s Office of National Marine Sanctuaries (ONMS) serves as the federal trustee for a system of National Marine Sanctuaries and Marine National Monuments in the ocean and Great Lakes that were established for resource protection, research, education, and sustainable public use. Sanctuary research and monitoring programs support management efforts by assessing sanctuary change. Conditions within a sanctuary are influenced by a wide variety of ocean and climate drivers, ecological processes, human activities, and actions of adjacent and overlapping management and regulatory entities. One reporting tool developed by ONMS, and first released in 2007, is the sanctuary Condition Report, which provides a summary of resources in a sanctuary, pressures on those resources, current conditions and trends, and management responses to various pressures on the sanctuary ecosystem (ONMS, 2018). These reports are structured around standardized questions that provide a tool for reporting on the status and trends of human pressures, water quality, habitat, living resources, maritime heri-

tage resources, and ecosystem services. Although the structure of a Condition Report is standardized nationally, the content of a report and the indicators used to address questions are sanctuary-specific; therefore, the completeness and impact of a Condition Report depends on the existence and availability of locally relevant observations and their synthesis.

Sanctuary Condition Reports are a fairly new management tool, and so far have been completed at seven- to 10-year intervals. A significant amount of time and resources are involved in the process of updating a report, a large part of which is compiling and summarizing the relevant monitoring data. For example, the first Condition Report for the MBNMS was published as a large document in 2009 (ONMS, 2009), and a partial update was published in 2015 (ONMS, 2015). Though these reports have provided extensive status and trend information at the time of publication, effective and timely resource management requires status and trend information that is readily updated as new data are collected. The information presented needs to be visually clear, publicly accessible, and understandable by both technical experts and the public.

IEA Program, ESRs, and Information Needs for Management

The need for ecosystem-based fisheries management has become heightened with a rapidly changing ocean and increased interactions between protected species and human activities. The California Current IEA (CCIEA) is managed in part through information from its Ecosystem Stats Reports (ESRs), which integrate ecosystem indicators from many sources including NOAA's Northwest and Southwest Fisheries Science Centers, fisheries landings, academia, and others (Samhoury et al., 2014; Levin et al., 2014; Harvey et al., 2019). These indicators have also been incorporated into risk analyses (Samhoury et al., 2017) and management strategy evaluations (Dawson and Levin, 2019). Analysis of data is simpli-

fied by using standard formats, including the Darwin Core standard for taxonomic data (Benson et al., 2021, in this issue), and serving them together in a web service (ERDDAP) that provides human and machine-to-machine access using software tools of the researcher's choice. Web services also enable interactive web page dashboards and infographics.

The IEA focuses broadly on ecosystem-based management, including human activities beyond living resource extraction, such as wind energy development, effects of marine traffic on ecosystems, and species range shifts or expansions across jurisdictional boundaries. The ESR is developed annually, focusing on northern, central, and southern regions of the California Current and reporting from physical oceanography up to human dimensions, and is presented to the Pacific Fisheries Management Council every March (Harvey et al., 2019). This report started as a ~1,000-page document (Levin and Schwing, 2011) and has evolved into a 20- to 30-page report with important appendices available online (NOAA, 2021a). Anomalous conditions and ecosystem perturbations have resulted in the appearance in the ESR of new sections focusing on the recent warm water event and human-wildlife conflict such as between whales and crab fisheries (Santora et al., 2020). The CCIEA is also moving toward increased automation and broader partnership beyond the council, recognizing that the tedium of incorporating new data annually limits the growth of new scientific aims.

DELIVERING BIODIVERSITY INFORMATION TO MEET MANAGEMENT REQUIREMENTS

A key challenge in developing a distributed information system is building a common understanding of priorities, requirements, actions, actors, structure, flow, and user values, as well as feedback to drive system improvement (e.g., Iwamoto et al., 2019). MBON helped shape workflows that led to the integration of biological observing information

for ocean-health indicator products (see also Benson et al., 2021, in this issue). The workflow facilitates the synthesis of data and integration into web-based tools that summarize ecosystem status and changing conditions. Across the nation, MBON successes include improved access to data repositories, technical innovation, and development of a suite of user-focused products that align with existing management requirements and are responsive to adaptive management needs (Figure 1).

The quality and availability of marine biodiversity observations are critical for developing products that adequately meet user needs (Bailey et al., 2019; Iwamoto et al., 2019). Building consensus for workflow elements and communicating them to the contributors and users via "datastream plans" leads to efficiency in information flow and delivery of products. Biological data needed for integrated assessments have been available from disparate individuals, networks, and teams, in a variety of formats and with sparse or poorly structured metadata (Buck et al., 2019). A collaborative network, with an operational focus in the IOOS Regional Associations is accelerating discovery and use of biological ocean data through the promotion and uptake of standardized data and metadata services and tools and the application of such workflow planning for access points such as infographics and dashboards.

Prioritizing Data Targets Based on Resource Management Requirements

From a management perspective, identifying and prioritizing observing data that align with policy requirements and management priorities is a critical first step (Table 1) toward achieving a workflow that maximizes management impact and focuses limited resource allocation (Figure 1). We assessed which existing and developing observation programs would satisfy multiple management requirements related to GOOS Essential Ocean Variables (EOVs), Sanctuary Condition Report questions (ONMS, 2018), and IEA program focal areas (Figure 1a, Table 1).

Monterey Bay National Marine Sanctuary priority indicators are regionally specific, align with one or more Condition Report questions, and inform conditions of one or more of the eight major habitats in MBNMS (Brown et al., 2019). Similarly, CCIEA ESR indicators have been selected to inform Pacific Fisheries Management Council needs (Harvey et al., 2019). This assessment process matched information needs from these management drivers to available regional observing data sources with particular attention to identifying data sources that filled multiple management needs.

The assessment identified several high-priority data sets for data integration and product development (Table 1), including remote-sensing satellite data (e.g., sea

surface temperature, chlorophyll-*a*), ship-based net samples (phytoplankton, zooplankton, forage fish, groundfish), shore-based observations (pinniped and seabird colonies, beached carcasses), and ship-based observations (seabirds, mammals, whale entanglements). Specific topics included indicators of biodiversity change at multiple trophic levels; local expression of, and impacts to, water quality and biodiversity during marine heatwaves; and causes of seabird and mammal mortality and whale entanglement. We also highlighted key observing platforms and methods of collecting data for top priority core variables and management requirements (Figure 1b).

Several technology advancements are also improving efficiency of collection

and delivery of management priority data sets about the ocean and its diverse species (e.g., Estes et al., 2021). Observations in the established areas of visual surveys, along with water, net, and sediment samples, join emerging methods such as plankton imaging, ocean sound, and multi-trophic level eDNA monitoring that are increasing the quantity and quality of information that supports dynamic management (Figure 1b). In the near-term, researchers and data scientists in the region will be collaborating via MBON and other initiatives to improve automation, including machine learning pipelines. These are steps that are taking what traditionally has been the realm of research into operational biological ocean observing.




FIGURE 1. Workflow organogram highlighting MBON-focused sequential and cyclical relationships among (a) policy, management, and prioritization of (b) observing platforms and methods. (c) Application to ocean observing practices and information handling. (d) The evolution of information provision for users in application-specific decision-making. Sections (c) and (d) embody FAIR (Findability, Accessibility, Interoperability, and Reusability) data principles (Wilkinson, 2016). Governance structures operate at various points throughout to implement each system component.

FAIR Data Handling and Services

A cornerstone of MBON is to expand capability and capacity for enduring biological and ecosystem data management, data stream planning, reliable dissemination through archives and portals, and the safeguarding of data through metadata and data formatting standards and archiving. An identified need is access to biodiversity data through public portals, such as the Ocean Biodiversity Information System (OBIS, 2021) or the new joint CeNCOOS/SCCOOS (Southern

California Coastal Ocean Observing System) Data Portal. This effort is guided by FAIR data principles (emphasizing Findability, Accessibility, Interoperability, and Reusability of digital assets by machines; Wilkinson et al., 2016). This is fundamental to contending with the rapidly increasing volume and complexity of data. In the future, these information delivery systems will be closely coupled with numerical models that will expand the fields spatially and forecast into the future. Models themselves will be part of

the quality control system.

Innovations in MBON, CCIEA, and IOOS data management systems include streamlined access to biology and ecosystems data (Benson et al., 2021, in this issue), enabled by data preparations that include quality control, the application of metadata standards, and data and archive services (Figure 1c). This work is supported by the Southwest Fisheries Science Center, CeNCOOS, MBON, and the IOOS Animal Telemetry Network (IOOS, 2021a), along with Earth Science

TABLE 1. Priority-setting matrix linking Global Ocean Observing System (GOOS) Essential Ocean Variables (EOVs) with Integrated Ecosystem Assessment (IEA) and sanctuary information needs and examples of observing data that meets those needs.

| GOOS EOVS | IEA FOCAL COMPONENT AND CCIEA ESR INDICATOR(S) | ONMS MANAGEMENT QUESTIONS | MBNMS-SPECIFIC INDICATOR(S) (Pelagic Habitat Focus) | REGIONAL OBSERVING DATA/ PRODUCT EXAMPLES |
|--|---|--|---|---|
| Phytoplankton biomass and diversity | | <ul style="list-style-type: none"> Question 12: Keystone and foundation species | <ul style="list-style-type: none"> Phytoplankton biomass/volume/taxonomic structure Chl-α | <ul style="list-style-type: none"> MBARI M1 phytoplankton counts Satellite-based Chl-α CalHABMAP and IFCBs |
| Zooplankton biomass and diversity | ECOLOGICAL INTEGRITY COMPONENT Forage availability | <ul style="list-style-type: none"> Question 12: Keystone and foundation species Question 15: Biodiversity | <ul style="list-style-type: none"> Krill and ichthyoplankton abundance and diversity | <ul style="list-style-type: none"> RREAS surveys ACCESS cruises |
| Invertebrate abundance and distribution | ECOLOGICAL INTEGRITY COMPONENT Forage availability, jellyfish biomass, trophic structure | <ul style="list-style-type: none"> Question 12: Keystone and foundation species Question 15: Biodiversity | <ul style="list-style-type: none"> Invertebrate abundance and diversity | <ul style="list-style-type: none"> RREAS surveys |
| Fish abundance and distribution | GROUND FISHES, SALMON, AND ECOLOGICAL INTEGRITY COMPONENTS Diversity, forage availability, trophic structure | <ul style="list-style-type: none"> Question 12: Keystone and foundation species Question 13: Other focal species Question 15: Biodiversity | <ul style="list-style-type: none"> Salmon abundance Forage fish abundance and diversity | <ul style="list-style-type: none"> RREAS surveys ATN data NWFSC groundfish surveys |
| Marine bird abundance and distribution | SEABIRD COMPONENTS At-sea population, mortality, productivity | <ul style="list-style-type: none"> Question 4: Human activities impacting living resources Question 13: Other focal species Question 15: Biodiversity | <ul style="list-style-type: none"> At-sea seabird abundance, distribution, biodiversity Nesting colony abundance Beached seabird counts | <ul style="list-style-type: none"> RREAS surveys BeachCOMBERS/Beach Watch |
| Marine mammal abundance and distribution | MARINE MAMMAL COMPONENTS Sea lion, cetacean; Ecological integrity | <ul style="list-style-type: none"> Question 4: Human activities impacting living resources Question 13: Other focal species | <ul style="list-style-type: none"> At-sea whale distribution and abundance Pinniped pup production Beached mammal counts Entangled whale counts | <ul style="list-style-type: none"> RREAS surveys NMFS MMTD surveys NMFS SMI pinniped surveys BeachCOMBERS/Beach Watch ATN data |
| Ocean sound | | <ul style="list-style-type: none"> Question 4: Human activities impacting living resources | <ul style="list-style-type: none"> Sound levels | <ul style="list-style-type: none"> Passive listening stations (e.g., MBARI hydrophone) |
| Sea surface and subsurface temperature | CLIMATE AND OCEAN DRIVERS Sea surface temperature | <ul style="list-style-type: none"> Question 8: Water conditions and climate change | <ul style="list-style-type: none"> Sea surface and seafloor temperature Air temperature | <ul style="list-style-type: none"> UCSC ROMS, Hopkins, and CeNCOOS sensors GHRSSST |

ACCESS = Applied California Current Ecosystem Studies; ATN = Animal Telemetry Network; CalHABMAP = California Harmful Algal Bloom Monitoring and Alert Program; CCIEA ESR = California Current Integrated Ecosystem Assessment Ecosystem Status Reports; CeNCOOS = Central and Northern California Observing System; GHRSSST = Group for High Resolution Sea Surface Temperature; IFCBs = Imaging FlowCytobots; MBARI = Monterey Bay Aquarium Research Institute; MBNMS = Monterey Bay National Marine Sanctuary; NMFS = National Marine Fisheries Service; MMTD = Marine Mammal and Turtle Division; NWFSC = Northwest Fisheries Science Center; ONMS = Office of National Marine Sanctuaries; RREAS = SWFSC Rockfish Recruitment and Ecosystem Assessment Survey; SMI = San Miguel Island; SWFSC = Southwest Fisheries Science Center; UCSC ROMS = University of California Santa Cruz Regional Ocean Modeling System

Information Partners (ESIP) programs. Examples of data preparation include the application of the Quality Assurance/Quality Control of Real Time Oceanographic Data (QARTOD; IOOS, 2021b) system to flag suspect or erroneous data, and (semi)automated processing of video and still-image photographic surveys, ocean sound, and marine genomic information including eDNA. Machine readable metadata standards including Ecological Metadata Language (EML) and Darwin Core (Wieczorek et al., 2012; TDWG, 2021) enable data on servers to feed into a variety of web portals and infographic tools. Metadata formatting using the OBIS ENV-DATA approach enables linking of a wide variety of information, including attributes like abundance, body size, and biomass, or habitat and environmental information related to the sample (DePooter et al., 2017). Practitioners follow international guidelines such as the GOA-ON metadata standard for biogeochemistry (Newton et al., 2015), the World Register of Marine Species (WoRMS Editorial Board, 2021), and those designed for the training of artificial intelligence tools.

The open source ERDDAP data server developed by NOAA provides a simple consistent way to make data available using a web interface or machine-to-machine transfer. This enables brokering between data centers with no separate dedicated infrastructure (Buck et al., 2019). Geospatial and model data are also frequently made available through Thematic Realtime Environmental Distributed Data Services (THREDDS) servers. For example, the Rockfish Recruitment and Ecosystem Assessment Survey (RREAS; Sakuma, et al., 2016; Santora et al., 2021, in this issue) was an early candidate for prototyping the MBON data flow process. The Fisheries Ecology Division (FED) of SWFSC has conducted this midwater trawl survey off central California since 1983 with the primary goal of developing pre-recruit indices for young-of-the-year (YOY) rockfish (*Sebastes* spp.). The survey also sam-

ples numerous other components of the epipelagic micronekton. These data were made available for machine-to-machine access via ERDDAP directly from a relational database (PostgreSQL), updated regularly, aligned to Darwin Core, uploaded to GBIF (Field et al., 2019), and made available in OBIS.

Co-development of Dynamic Data Products for Status and Trends

Delivering effective products for management and policy applications requires early and repeated coordination among data providers, product developers, and managers and other users, to facilitate collaboration and iterative product development (Figure 1d). The MBON team developed a number of products in collaboration with and for specific users, including sanctuary managers, scientists, educators, fisheries management councils, and CeNCOOS stakeholders. Some examples follow.

Interactive Infographics

The MBON project developed and implemented interactive infographics in collaboration with various groups, including the regional IEA programs, IOOS, and sanctuary teams. Interactive infographics deliver sanctuary-specific data products for assessing and interpreting ecosystem status and trends. The infographics use web-enabled technologies, including open-source software and data, to link illustrations of ecosystem elements to complex underlying time series of observations. This link allows users to quickly see trends and background information in a multimedia format. Interactive infographics have gained interest regionally, nationally, and internationally as a highly successful and innovative way to serve data visualizations.

For the California set of MBON efforts, the ecosystem illustrations were created in collaboration with the California Current IEA graphic designer (Su Kim, NOAA). They show Condition Report indicators in each of a sanctuary's major habitats (background scene) using eco-

system elements (e.g., icons of fish, plankton, kelp, and climate drivers and human uses) (Figure 2a; Brown et al., 2019; Spector et al., 2021). An interactive infographic framework was developed (using D3 JavaScript, R, and Rmarkdown software) to link the ecosystem elements in the illustration to pop-up windows showing one or more figures along with the latest associated data visualization (e.g., time series, map, chart). The websites and versioned source code are hosted at <https://github.com/> and updated using the Travis continuous integration web service. The storage requirements are modest (<1 GB) and computation requirements are also minimal (<1 hr). Example MBON infographic products are available on GitHub (Best and Ranganathan, 2021). Infographics code is free, open for use by anyone, and described in more detail in Spector et al. (2021).

MBON and CeNCOOS Portals with Curated Data Views

Sanctuary science teams and advisory panels may on occasion need access to research and assessment tools with more technical capabilities than are offered in the interactive infographics. For more expert users, the ability to discover, explore, and manipulate data is available through the MBON and the new joint CeNCOOS/SCCOOS Data Portals (Figure 2c) and catalog tools.

Customizable curated data views such as those generated within these portals show multiple graphical plots that relate to a specific scientific or Condition Report question or to a management issue (Figure 2b). Metrics and graphs are predetermined to ensure that all users are viewing and evaluating the same information, but the user can change time-scale and other features to explore the displayed information. These views can illustrate change over time and space for specific areas and analytics that describe relationships between variables, and they provide context for change in larger regions or reference areas. For example, one view might illustrate patterns

in satellite-based sea surface temperature and species richness of the pelagic forage community, along with encounter rates of seabird carcasses. Curated data views can be modified to update the data for every viewing or on a periodic basis, and the URL can be copied and sent to other users, who could then render the same catalog view and illustrations on a separate computer.

Options for curated data views include satellite, ocean model, and in situ data sets from large programs, smaller initiatives, and individual researchers and citizen scientists. Tools available in the CeNCOOS and MBON portals have been developed that quantify changes in regional climate variation, sea surface temperature, heat content, surface currents, wave height and direction, and net primary production, and these can be combined with observations of habitat, biomass, community composition and/

or species change. Summary forms of this information can be provided as ecological indicators and report card categorizations of conditions. Analytics for specific marine spaces and reference areas can also be digested for inclusion into infographics and other tools suitable for more a general audience.

Maturing Technologies and Data Streams for Biology and Biodiversity Observing

The policy, management, and related observing drivers identified in Table 1 motivate the development of new methods and their transitions from research into sustained applied observations. Biological observing methods that can address priority needs and that are becoming more feasible include those in genomics, imaging, animal tagging, and passive acoustic and satellite-based monitoring (see also Estes et al., 2021).

Environmental DNA (eDNA)

Traditional field methods for observing biodiversity are expensive, require intensive human intervention, and are time-consuming. Examples include ship expeditions that use net trawls or visual diver surveys in shallower waters. In most cases, these surveys only cover one or two trophic levels, a limited number of taxa, and rely on expert taxonomists. Environmental DNA is a method to survey organisms that may be present in an area by examining the genetic material left behind suspended in the seawater. The approach is akin to that of a forensic scientist who searches for human material left behind at a crime scene, extracts the DNA from that material, and compares it to a bank of suspect DNA to find a match. In the case of eDNA, the material left behind by the suspects is collected by filtration and our suspects are life in the sea at every trophic level.




FIGURE 2. Tiered data products targeting different audiences from general to expert users. (a) Interactive infographics provide status and trend information that is visually appealing, publicly accessible, and easily understood by both technical and nontechnical users. (b) Curated data views provide updates to a set of interrelated metrics for a more involved user, such as those involved with a monitoring program, a management issue, or an advisory group. (c) The MBON Data Portal and new joint Central and Northern California Ocean Observing System/Southern California Coastal Ocean Observing System Data Portal facilitate data visualization, exploration, and manipulation by the most technically adept users, including data providers and subject experts.

CASE STUDY

BeachCOMBERS: A California Network of Volunteer Observers

A case study from the BeachCOMBERS seabird and mammal stranding survey initiative provides an example of prioritization to end-user workflow (Figure 1). When a stranding or mortality event is observed on local beaches, it can be challenging to determine whether the event is natural or related to issues such as harmful algal blooms (HABs), oil spills and seeps, ingestion and entanglement in marine debris and fishing gear, local prey availability, or vessel strikes. Detailed information on extent and duration of wild-life mortality events can help managers determine causes and develop mitigation strategies for the future. In the fall of 2015, increased numbers of dead and stranded juvenile common murrelets were observed on central California beaches. This was later linked to the prevalence of warm water and low prey conditions along the coast (Gibble et al., 2018) and then to an extended marine heatwave (Piatt et al., 2020). CeNCOOS worked with data providers to aggregate beach surveys of marine mammals and seabirds from BeachCOMBERS, Beach Watch citizen science programs, and the Marine Mammal Stranding Program at Humboldt State University. These data now form the “Effort-based surveys, Northern and Central California Beaches: Seabirds and Marine Mammals” data product that combines surveys over a 25-year period from Del Norte to Los Angeles Counties. This harmonized data product is now available for exploring through the CeNCOOS/SCCOOS Data Portal.

Once data are available through ERDDAP data services, tiered data products are developed with input from BeachCOMBERS and sanctuary resource protection and education staff (Figure 2). Integration of seabird and mammal stranding data into the CeNCOOS/SCCOOS portal allows BeachCOMBERS researchers and partners to easily visualize observations to better explore potential causes of seabird and marine mammal strandings and inform potential management actions to reduce wildlife impacts from human activities. To engage a wider audience that includes managers, educators, and the interested public, we incorporated BeachCOMBERS summary data into interactive infographics as described above (Figure 2a). A set of curated data views were developed to showcase the same metrics regularly provided by BeachCOMBERS in their periodic technical reports (Figure 2b) and used in sanctuary Condition Reports. The most technically adept users of MBON data products, including data providers and subject experts, will be able to discover, explore, and manipulate data using the CeNCOOS/SCCOOS portal (Figure 2c). Lastly, seabird mortality rates based on stranded carcass counts from BeachCOMBERS and Beach Watch are presented as indicators of seabird mortality rates in the CCIEA’s annual Ecosystem Status Report to the PFM (Harvey et al., 2019) and in the CCIEA (NOAA, 2021a,b).

During the first Sanctuary MBON project, the team developed best practices for using eDNA including the use of a routine set of markers that cover the tree of life, from microbes to whales (Chavez et al., 2021, in this issue). The team made systematic collections throughout the Florida Keys and Monterey Bay National Marine Sanctuaries, as well as offshore Santa Barbara, Hawai’i, and Peru. Frozen samples were exploited to develop decade-long time series. The results, together with many others in the growing literature, show that eDNA is a useful method for discriminating changes in marine communities over time and space (horizontally from sanctuary to sanctuary and vertically with depth). During the follow-on MBON projects, the teams in different projects are working on resolving many of the challenges associated with eDNA as well as exploiting the opportunities. Sampling is being expanded to the full CeNCOOS domain and elsewhere, and efforts are being made to transition the work from research to operations, such as in data processing and management to deliver information on the occurrences of managed and protected species.

Phytoplankton Surveys and HAB Monitoring

Biological sampling of phytoplankton is frequently limited by logistical constraints (including geographic location, depth, sample acquisition methodology, handling, processing). As a result, historical sampling density is considerably more coarse spatially and temporally than physical and chemical measurements. The rapid maturity of new technologies such as the Imaging FlowCytobot (IFCB) begins to address these issues. The IFCB is an automated flow cytometer with imaging capabilities (i.e., an automated microscope; Sosik and Olson, 2007). It can sample every 25 minutes from fixed locations, ships, and autonomous vehicles. When the IFCB is coupled with tools for automated classification of images, the system produces phytoplankton abundance estimates of species- and genus-level taxa in near-real time (e.g., Fischer et al., 2020). Four instruments have routinely been deployed off California with support from NOAA and the National Science Foundation. In 2020, the California Ocean Protection Council expanded this network to include 10 instruments, with two deployed on coastal moorings and several alternating between shipboard and fixed location sampling. One goal is to support the California Harmful Algal Bloom Monitoring and Alert Program (HABMAP). The network will also be used to explore the largely beneficial plankton community structure and biodiversity across a wide swath of the California Current System, as this is the food and energy foundation for the coastal marine ecosystem in this region.

A key metric for success is the establishment of coordinated operations and centralized data management designed to scale to a national network as such data become increasingly common within IOOS. For example, the rich data set from the Santa Cruz Municipal Wharf (Figure 3) is becoming more widely available. The HAB organism *Pseudo-nitzschia* is quickly and reliably quantified using the IFCB and computer vision tools. The data also show an unusual dominance of dinoflagellates in 2018 related to local environmental forcing (Fischer et al., 2020). The true power of these analyses will be realized when the full suite of 10 instruments can provide a picture of plankton dynamics across the California Current system, including HAB species bloom extent and duration.

Imaging with Cameras on Gliders and Other Platforms

The recent development of a microscopic camera system for gliders is paving the way to greater availability of marine snow and zooplankton abundance and diversity information (Lombard et al., 2019). For example, the Zooglider, developed at the Scripps institution of Oceanography, has photographic and acoustic zooplankton sensing capability (Ohman et al., 2019). Image data can be processed using automated object detection and classification. Expanded use of such tools on gliders

and long-range autonomous underwater vehicles in IOOS could provide profiles and repeat ocean transects of zooplankton variables with high spatial and temporal resolution. This could deliver data for the Ecological Integrity category of the CCIEA, specifically copepods, including northern and southern copepods as well as krill including *Euphausia*, *Pyrosoma*, *Aurelia*, *Chrysaora*, *Aequorea*, and fish larvae. These tools could also deliver biodiversity information for MBON and the MBNMS “Zooplankton species/abundances” Condition Report indicator. Other tools are also maturing using a variety of approaches including holography, structured light, and confocal imaging across a wide range of object sizes (Lombard et al., 2019).

Animal Tagging

The Animal Telemetry Network (ATN) is building partnerships and data handling capability for animal tag sensor data. This method can collect information about the ecology and physiology of tagged individuals and serve as a source of operational oceanographic data (Sequeira et al., 2018; Harcourt et al., 2019). Depending on the tagged species, thousands of profiles can be collected on temperature, salinity, and other information to hundreds of meters depth and/or ocean sections covering thousands of kilometers in range (e.g., Naito et al., 2017; McHuron et al.,

2018). By tagging across several taxa, such as tuna, sharks, and seals, it is possible to cover a wide range of ocean observations that can provide critical information for how changes in the MBNMS interact with variations occurring in the greater northeastern Pacific. Critically, this information can come associated with the movements of managed species and has policy applications such as understanding ranges, connectivity, and movements in and out of marine protected areas (Dunn et al., 2019).

Ocean Soundscapes

Passive acoustic monitoring (PAM) is essential to understanding marine biodiversity (Mooney et al., 2020). By enabling detection of the presence and behavior of soniferous animals, including mammals and fish, PAM expands knowledge of the species inhabiting an ecosystem far beyond what can be achieved using only visual surveys. Because sound travels so effectively in marine waters, PAM can also provide far-reaching detection ranges, up to hundreds of kilometers for the powerful, low-frequency calls of endangered blue and fin whales (Figure 4). Further, the temporal coverage of biodiversity monitoring for soniferous species is greatly enhanced by the potential for PAM to provide continuous data. Integrative application of PAM and animal-borne sensing in MBNMS




FIGURE 3. Daily time series of phytoplankton-specific carbon fractions (top) and *Pseudo-nitzschia* spp. cell concentrations (bottom) generated by an Imaging FlowCytobot (IFCB) at the Santa Cruz Municipal Wharf (Monterey Bay, California) in 2018 and a machine learning image classifier. The top panel shows the fraction of phytoplankton-specific carbon composed of dinoflagellate taxa (pinks, reds, oranges, yellows), diatom taxa (greens, blues, purples), and nanoplankton (gray). The bottom panel shows the same time series for the toxic diatom genus *Pseudo-nitzschia* compared to weekly Fluorescent In Situ Hybridization (FISH) microscopy of discrete samples; the inset image provides an example of an IFCB image for *Pseudo-nitzschia*. White bands indicate data gaps. Adapted from Fischer et al. (2020)

has enabled detection of blue whale migration, a new capability applicable to protection of this endangered species (Oestreich et al., 2020). Together with effective ecosystem monitoring, relationships between ecosystem dynamics and marine mammal behavior can be examined (Ryan et al., 2019). This method is now being tested for long-term monitoring in marine sanctuaries (SanctSound program) and in the context of marine mammal indicator data for IEAs.

By enabling detection of anthropogenic noise in the ocean, PAM also supports research that is essential to an emergent dimension of ocean stewardship—protection of acoustic habitat (Southall et al., 2009; Gedamke et al., 2016; Hatch et al., 2016). While acute noise sources have received attention due to dramatic impacts on marine mammals observed in some regions, chronic noise is recognized as a global issue that must be addressed regionally through ecosystem-based management (Hatch et al., 2016). Within MBNMS, noise from seal bombs—explosives used in fishing operations with the goal of deterring interference in fishing operations by sea lions, seals, and dolphins—has been a focal research topic (Simonis et al., 2020). PAM revealed occurrence patterns, with up to

88 explosions per hour, while modeling of acoustic transmission revealed the far reach of noise from individual explosions across tens of kilometers. Persistent PAM in MBNMS has also emphasized the ephemeral nature of noise pollution and the associated potential for rapid improvement in habitat quality through noise reduction, as diminished shipping traffic during the COVID-19 pandemic significantly reduced noise within the frequency band used by baleen whales to communicate (Ryan et al., 2021).

ACHIEVING THE VISION OF MBON

Marine ecosystems are undergoing rapid change (Bindoff et al., 2019; Blowes et al., 2019). Around the globe, rates of change in species distribution, abundance, and biodiversity often outpace our ability to monitor or measure them. The CCLME harbors a diverse array of life, including some of the most diverse algal flora and marine mammal fauna in the world (Checkley and Barth, 2009; Fautin et al., 2010; Block et al., 2011). Its resources sustain major economic activity and food security. The CCLME is also one of the most robustly observed systems in the world (e.g., Chavez et al., 2017; Barth, 2018; Chan et al., 2019; Gallo

et al., 2019). Yet even in this well-studied region, major challenges persist in collecting and delivering biological and ecosystem data and information suitable for understanding, managing, and adapting to ecosystem changes and human activities. Anthropogenic pressures include climate change and ocean acidification, as well as the tourism, shipping, and natural resource extraction industries.

As data management and information dissemination tools evolve in our region and across the IOOS Regional Associations, the capacity to deliver more MBON products with (semi-) automated updating can become a standard for sanctuaries and IEAs across the national system. Biological data at its most advanced or “mature” stage is defined by the generation of routinely available and relevant information products (IOOC BIO-TT, 2016). Observational data collection approaches, including those that are currently maturing from research to operational domains, are delivering a greater selection of the EOVs in biology and ecosystems. This, in turn, will empower an evolution in estimating and modeling the current ecological state, and forecasting into the future, and it will require improved quantity and quality of data in order to parameterize models and improve them through assimilation. The processes described here are now also benefiting condition report work for the Cordell Bank and Greater Farallones National Marine Sanctuaries, as well as marine protected area assessment in California.

Ecological forecasting examples from the central and northern California region include EcoCast, which guides fishing based on drivers of both target and bycatch distribution estimation (Hazen et al., 2018); the California-Harmful Algal Bloom Risk Mapping (C-HARM) model (Anderson et al., 2016); and physical, biogeochemical, and ecological modeling of the California Current that includes an iteration of the North Pacific Ecosystem Model for Understanding Regional Oceanography




FIGURE 4. Acoustic indices of blue and fin whale vocal activity recorded in Monterey Bay National Marine Sanctuary through the Monterey Accelerated Research System (MARS) cabled observatory. The index is a calibrated signal-to-noise ratio: signal from whale song (rhythmic repeated sequences of sound produced by males of each species) and noise from background (adjacent frequency bands). For each month, the interquartile range and median are shown. *Whale artist: Larry Foster*

(NEMURO; Kishi et al., 2007; Fiechter et al., 2014). There are excellent opportunities for using and improving these tools more effectively and for including such capabilities in anticipated advancements in modeling such as the use of the West Coast Operational (Ocean) Forecast System (Kurapov et al., 2017) and addition of biogeochemical and ecological modeling capabilities.

Integration of biology, biodiversity, and ecosystem observations into a coherent system of environmental observing could bring valuable information to new and diverse users. Coordination is a limiting factor in the ability of researchers and funding entities to adequately serve stakeholder communities and the public. In order to deliver information at the level of the CCLME, better synchronization of information resources could be achieved through a “West Coast Marine Biodiversity Observing Network” encompassing the West Coast Ocean Alliance, the West Coast IOOS Regional Associations, and ONMS and IEA efforts, as well as the many governmental, non-governmental, academic, industry, education, and other organizations that make up these associations. This West Coast synchronization could build consensus and improve efficiencies among researchers and observing practitioners, help to coordinate funding resources, and promote a more effective understanding of our marine biological resources and ecosystems. Moreover, such an effort can inform possible practices for mitigation or adaptation to change, including expansion in the renewable energy, desalination, and aquaculture industries. Implementing these practices will require prioritizing increased knowledge of the marine environment and the life it supports. 🌐

REFERENCES

- Anderson, C.R., R.M. Kudela, M. Kahru, Y. Chao, L.K. Rosenfeld, F.L. Bahr, D.M. Anderson, and T.A. Norris. 2016. Initial skill assessment of the California Harmful Algae Risk Mapping (C-HARM) system. *Harmful Algae* 59:1–18, <https://doi.org/10.1016/j.hal.2016.08.006>.
- Bailey, K., C. Steinberg, C. Davies, G. Galibert, M. Hidas, M.A. McManus, T. Murphy, J. Newton, M. Roughan, and A. Schaeffer. 2019. Coastal monitoring observing networks and their data products: Recommendations for the next decade. *Frontiers in Marine Science* 6:180, <https://doi.org/10.3389/fmars.2019.00180>.
- Barth, J.A., J.P. Fram, E.P. Dever, C.M. Risien, C.E. Wingard, R.W. Collier, and T.D. Kearney. 2018. Warm blobs, low-oxygen events, and an eclipse: The Ocean Observatories Initiative Endurance Array captures them all. *Oceanography* 31:90–97, <https://doi.org/10.5670/oceanog.2018.114>.
- Bax, N.J., P. Miloslavich, F.E. Muller-Karge, V. Allain, W. Appeltans, S.D. Batten, L. Benedetti-Cecchi, P.L. Buttigieg, S. Chiba, D.P. Costa, and others. 2019. A response to scientific and societal needs for marine biological observations. *Frontiers in Marine Science* 6:395, <https://doi.org/10.3389/fmars.2019.00395>.
- Benson, A., T. Murray, G. Canonico, E. Montes, F.E. Muller-Karger, M.T. Kavanaugh, J. Triñanes, and L.M. deWitt. 2021. Data management and interactive visualizations for the evolving Marine Biodiversity Observation Network. *Oceanography* 34(2):130–141, <https://doi.org/10.5670/oceanog.2021.220>.
- Best, B., and J. Ranganathan. 2021. “Guide to Infographiq,” version 2021-10-07, <https://marinebon.org/infographiq/>.
- Bindoff, N.L., W.W.L. Cheung, J.G. Kairo, J. Aristegui, V.A. Guinder, R. Hallberg, N. Hilmi, N. Jiao, M.S. Karim, L. Levin, and others. 2019. Changing ocean, marine ecosystems, and dependent communities. Pp. 447–587 in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem, J. Petzold, B. Rama, and N.M. Weyer, eds.
- Block, B.A., I.D. Jonsen, S.J. Jorgensen, A.J. Winship, S.A. Shaffer, S.J. Bograd, E.L. Hazen, D.G. Foley, G.A. Breed, A.L. Harrison, and J.E. Ganong. 2011. Tracking apex marine predator movements in a dynamic ocean. *Nature* 475(7354):86–90, <https://doi.org/10.1038/nature10082>.
- Blowes, S.A. 2019. The geography of biodiversity change in marine and terrestrial assemblages. *Science* 366:339–345, <https://doi.org/10.1126/science.aaw1620>.
- Bond, N.A., M.F. Cronin, H. Freeland, and N. Mantua. 2015. Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophysical Research Letters* 42:3,414–3,420, <https://doi.org/10.1002/2015GL063306>.
- Brown, J., G.D. Williams, C.J. Harvey, A.D. DeVogelaere, and C. Caldwell. 2019. *Developing Science-Based Indicator Portfolios for National Marine Sanctuary Condition Reports*. Marine Sanctuaries Conservation Series ONMS-19-07, US Department of Commerce, National Oceanic and Atmospheric Administration, Office of National Marine Sanctuaries, Silver Spring, MD, 78 pp., <https://sanctuaries.noaa.gov/science/conservation/2019-science-based-indicator-portfolios.html>.
- Buck, J.J.H., S.J. Bainbridge, E.F. Burger, A.C. Kraberg, M. Casari, K.S. Casey, L. Darroch, J. Del Rio, K. Metfies, E. Delory, and others. 2019. Ocean data product integration through innovation: The next level of data interoperability. *Frontiers in Marine Science* 6:32, <https://doi.org/10.3389/fmars.2019.00032>.
- CeNCOOS (Central and Northern California Ocean Observing System). 2020. *CeNCOOS Strategic Plan (2020-25): Advancing Ocean Observing in Central & Northern California*. CeNCOOS, Moss Landing, CA, 34 pp., <https://www.cencoos.org/images/docs/CeNCOOS-Strat-Plan-2020.pdf>.
- Chan, F., J.A. Barth, K.J. Kroeker, J. Lubchenco, and B.A. Menge. 2019. The dynamics and impact of ocean acidification and hypoxia: Insights from sustained investigations in the Northern California Current Large Marine Ecosystem. *Oceanography* 32(3):62–71, <https://doi.org/10.5670/oceanog.2019.312>.
- Chavez, F.P., J. Ryan, S.E. Lluch-Cota, and M. Niquen. 2003. From anchovies to sardines and back: Multidecadal change in the Pacific Ocean. *Science* 299(5604):217–221, <https://doi.org/10.1126/science.1075880>.
- Chavez, F.P., and M. Messie. 2009. A comparison of eastern boundary upwelling ecosystems. *Progress in Oceanography* 83:80–96, <https://doi.org/10.1016/j.pocean.2009.07.032>.
- Chavez, F.P., J.T. Pennington, R.P. Michisaki, M. Blum, G.M. Chavez, J. Friederich, B. Jones, R. Herlien, B. Kieft, B. Hobson, and others. 2017. Climate variability and change: Response of a coastal ocean ecosystem. *Oceanography* 30(4):128–145, <https://doi.org/10.5670/oceanog.2017.429>.
- Chavez, F.P., M. Min, K. Pitz, N. Truelove, J. Baker, D. LaScala-Grunewald, M. Blum, K. Walz, C. Nye, A. Djurhuus, and others. 2021. Observing life in the sea using environmental DNA. *Oceanography* 34(2):102–119, <https://doi.org/10.5670/oceanog.2021.218>.
- Checkley, D.M. Jr., and J.A. Barth. 2009. Patterns and processes in the California Current System. *Progress in Oceanography* 83(1–4):49–64, <https://doi.org/10.1016/j.pocean.2009.07.028>.
- Dawson, C., and P.S. Levin. 2019. Moving the ecosystem-based fisheries management mountain begins by shifting small stones: A critical analysis of EBFM on the US West Coast. *Marine Policy* 100:58–65, <https://doi.org/10.1016/j.marpol.2018.11.005>.
- De Pooter, D., W. Appeltans, N. Bailly, S. Bristol, K. Deneudt, M. Eliezar, E. Fujioka, A. Giorgetti, P. Goldstein, M. Lewis, and others. 2017. Toward a new data standard for combined marine biological and environmental datasets – Expanding OBIS beyond species occurrences. *Biodiversity Data Journal* 5:e10989, <https://doi.org/10.3897/BDJ.5.e10989>.
- Di Lorenzo, E., N. Schneider, K.M. Cobb, P.J.S. Franks, K. Chhak, A.J. Miller, J.C. McWilliams, S.J. Bograd, H. Arango, E. Curchitser, and T.M. Powell. 2008. North Pacific Gyre Oscillation links ocean climate and ecosystem change. *Geophysical Research Letters* 35(8), <https://doi.org/10.1029/2007GL032838>.
- Dunn, D.C., A.L. Harrison, C. Curtice, S. DeLand, B. Donnelly, E. Fujioka, E. Heywood, C.Y. Kot, S. Poulin, M. Whitten, and others. 2019. The importance of migratory connectivity for global ocean policy. *Proceedings of the Royal Society B* 286:20191472, <https://doi.org/10.1098/rspb.2019.1472>.
- Estes, M., C. Anderson, W. Appeltans, N. Bax, N. Bednaršek, G. Canonico, S. Djavidnia, E. Escobar, P. Fietzek, M. Gregoire, and others. 2021. Enhanced monitoring of life in the sea is a critical component of conservation management and sustainable economic growth. *Marine Policy* 132:104699, <https://doi.org/10.1016/j.marpol.2021.104699>.
- Fautin, D., P. Dalton, L.S. Incze, J.-A.C. Leong, C. Pautzke, A. Rosenberg, P. Sandifer, G. Sedberry, J.W. Tunnell Jr., I. Abbott, and others. 2010. An overview of marine biodiversity in United States waters. *PLoS ONE* 5:e11914, <https://doi.org/10.1371/journal.pone.0011914>.
- Fiechter, J., E.N. Curchitser, C.A. Edwards, F. Chai, N.L. Goebel, and F.P. Chavez. 2014. Air-sea CO₂ fluxes in the California Current: Impacts of model resolution and coastal topography. *Global Biogeochemical Cycles* 28:371–385, <https://doi.org/10.1002/2013GB004683>.

- Field, J., K. Sakuma, A. Benson, and L. deWitt. 2019. Rockfish Recruitment and Ecosystem Assessment Survey, Catch Data. Version 1.3, United States Geological Survey, Sampling event dataset, <https://doi.org/10.15468/73tvnd>, accessed via GBIF.org on 2020-08-12.
- Fischer, A.D., K. Hayashi, A. McGaraghan, and R.M. Kudela. 2020. Return of the “age of dinoflagellates” in Monterey Bay: Drivers of dinoflagellate dominance examined using automated imaging flow cytometry and long-term time series analysis. *Limnology and Oceanography* 65:2,125–2,141, <https://doi.org/10.1002/lno.11443>.
- Gallo, N.D., E. Drenkard, A.R. Thompson, E.D. Weber, D. Wilson-Vandenberg, S. McClatchie, J.A. Koslow, and B.X. Semmens. 2019. Bridging from monitoring to solutions-based thinking: Lessons from CalCOFI for understanding and adapting to marine climate change impacts. *Frontiers in Marine Science* 6:695, <https://doi.org/10.3389/fmars.2019.00695>.
- García-Reyes, M., and W.J. Sydeman. 2017. California Multivariate Ocean Climate Indicator (MOCI) and marine ecosystem dynamics. *Ecological Indicators* 72:521–529, <https://doi.org/10.1016/j.ecolind.2016.08.045>.
- Gedamke, J., J. Harrison, L. Hatch, R. Angliss, J. Barlow, C.L. Berchok, C. Caldwell, M. Castellote, D. Cholewiak, M.L. DeAngelis, and others. 2016. NOAA Ocean Noise Strategy Roadmap. National Oceanic and Atmospheric Administration, Silver Spring, MD, 144 pp., <https://cetsound.noaa.gov/road-map>.
- Gibble, C., R. Duerr, B. Bodenstein, K. Lindquist, J. Lindsey, J. Beck, L. Henkel, J. Roletto, J. Harvey, and R. Kudela. 2018. Investigation of a large-scale common murre (*Uria aalge*) mortality event in California, USA, in 2015. *Journal of Wildlife Diseases* 54:569–574, <https://doi.org/10.7589/2017-07-179>.
- Harcourt, R., A.M.M. Sequeira, X. Zhang, F. Roquet, K. Komatsu, M. Heupel, C. McMahon, F. Whoriskey, M. Meekan, G. Carroll, and others. 2019. Animal-borne telemetry: An integral component of the ocean observing toolkit. *Frontiers in Marine Science* 6:326, <https://doi.org/10.3389/fmars.2019.00326>.
- Harvey, C.J., N. Garfield, G.D. Williams, N. Tolimieri, I. Schroeder, K.S. Andrews, K. Barnas, E. Bjorkstedt, S. Brograd, R. Brodeur, and others. 2019. *Ecosystem Status Report of the California Current for 2019: A Summary of Ecosystem Indicators Compiled by the California Current Integrated Ecosystem Assessment Team (CCIEA)*. US Department of Commerce, NOAA Technical Memorandum NMFS-NWFSC-149, 101 pp., <https://doi.org/10.25923/p0ed-ke21>.
- Harvey, C.J., D.L. Fluharty, M.J. Fogarty, P.S. Levin, S.A. Murawski, F.B. Schwing, R.L. Shuford, C.R. Kelble, and M.E. Monaco. 2020. The origin of NOAA's Integrated Ecosystem Assessment Program: A retrospective and prospective. *Coastal Management* 49:9–25, <https://doi.org/10.1080/08920753.2021.1846110>.
- Hatch, L.T., C.M. Wahle, J. Gedamke, J. Harrison, B. Laws, S.E. Moore, J.H. Stadler, and S.M. Van Parijs. 2016. Can you hear me here? Managing acoustic habitat in US waters. *Endangered Species Research* 30:171–186, <https://doi.org/10.3354/esr00722>.
- Hazen, E.L., K.L. Scales, S.M. Maxwell, D.K. Briscoe, H. Welch, S.J. Bograd, H. Bailey, S.R. Benson, T. Eguchi, H. Dewar, S. Kohin, and others. 2018. A dynamic ocean management tool to reduce bycatch and support sustainable fisheries. *Science Advances* 4(5):3001, <https://doi.org/10.1126/sciadv.aar3001>.
- Hildreth, R. 2008. Place-based ocean management: Emerging U.S. law and practice. *Ocean & Coastal Management* 51:659–670, <https://doi.org/10.1016/j.ocecoaman.2008.07.005>.
- IOOC BIO-TT. (IOOC Biological Integration and Observation Task Team). 2016. *Biological and Ecosystem Observations Within United States Waters II: A Workshop Report to Inform Priorities for the United States Integrated Ocean Observing System*. National Ocean Council, ed., Washington, DC, http://www.iooc.us/wp-content/uploads/biological_and_ecosystem_observations_within_united_states_waters2.pdf.
- IOOS (Integrated Ocean Observing System). 2021a. Animal Telemetry Network, <https://atn.ioos.us/>.
- IOOS. 2021b. Quality Assurance/Quality Control of Real Time Oceanographic Data, <https://ioos.noaa.gov/project/qartod/>.
- Iwamoto, M.M., J. Dorton, J. Newton, M. Yerta, J. Gibeau, T. Shyba, B. Kirkpatrick, and R. Currier. 2019. Meeting regional, coastal and ocean user needs with tailored data products: A stakeholder-driven process. *Frontiers in Marine Science* 6:290, <https://doi.org/10.3389/fmars.2019.00290>.
- Kavanaugh, M.T., T. Bell, D. Catlett, M.A. Cimino, S.C. Doney, W. Klajbor, M. Messié, E. Montes, F.E. Muller-Karger, D. Otis, and others. 2021. Satellite remote sensing and the Marine Biodiversity Observation Network: Current science and future steps. *Oceanography* 34(2):62–79, <https://doi.org/10.5670/oceanog.2021.215>.
- Kishi, M.J., M. Kashiwai, D.M. Ware, B.A. Megrey, D.L. Eslinger, F.E. Werner, M. Noguchi-Aita, T. Azumaya, M. Fujii, S. Hashimoto, and others. 2007. NEMURO—a lower trophic level model for the North Pacific marine ecosystem. *Ecological Modelling* 202(1–2):12–25, <https://doi.org/10.1016/j.ecolmodel.2006.08.021>.
- Kurapov, A.L., S.Y. Erofeeva, and E. Myers. 2017. Coastal sea level variability in the US West Coast Ocean Forecast System (WCOSF). *Ocean Dynamics* 67:23, <https://doi.org/10.1007/s10236-016-1013-4>.
- Levin, P.S., and F. Schwing. 2011. Technical background for an integrated ecosystem assessment of the California Current: Groundfish, salmon, green sturgeon, and ecosystem health. US Department of Commerce, NOAA Technical Memorandum NMFS-NWFSC-109, 330 pp., <https://repository.library.noaa.gov/view/noaa/3842>.
- Levin, P.S., C.R. Kelble, R.L. Shuford, C. Ainsworth, Y. deReynier, R. Dunsmore, M.J. Fogarty, K. Holman, E.A. Howell, M.E. Monaco, S.A. Oakes, and F. Werner. 2014. Guidance for implementation of integrated ecosystem assessments: A US perspective. *ICES Journal of Marine Science* 71:1,198–1,204, <https://doi.org/10.1093/icesjms/fst112>.
- Lombard, F., E. Boss, A.M. Waite, M. Vogt, J. Uitz, L. Stemann, H.M. Sosik, J. Schulz, J.-B. Romagnan, M. Picherl, and others. 2019. Globally consistent quantitative observations of planktonic ecosystems. *Frontiers in Marine Science* 6:196, <https://doi.org/10.3389/fmars.2019.00196>.
- Mantua, N.J., S.R. Hare, Y. Zhang, J.M. Wallace, and R.C. Francis. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society* 78(6):1,069–1,080, [https://doi.org/10.1175/1520-0477\(1997\)078<1069:APICOW>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<1069:APICOW>2.0.CO;2).
- McHuron, E.A., B.A. Block, and D.P. Costa. 2018. Movements and dive behavior of juvenile California sea lions from Año Nuevo Island. *Marine Mammal Science* 34:238–249, <https://doi.org/10.1111/mms.12449>.
- Monaco, M.E., E. Spooner, S.A. Oakes, C.J. Harvey, and C.R. Kelble. 2021. Introduction to the NOAA Integrated Ecosystem Assessment Program: Advancing ecosystem based management. *Coastal Management* 49:1–8, <https://doi.org/10.1080/08920753.2021.1846109>.
- Mooney, T.A., L. Di Iorio, M. Lammers, T.-H. Lin, S.L. Nedelec, M. Parsons, C. Radford, E. Urban, and J. Stanley. 2020. Listening forward: Approaching marine biodiversity assessments using acoustic methods. *Royal Society Open Science* 7:201287, <https://doi.org/10.1098/rsos.201287>.
- Muller-Karger, F.E., P. Miloslavich, N.J. Bax, S. Simmons, M.J. Costello, I. Sousan Pinto, G. Canonico, W. Turner, M. Gill, E. Montes, and others. 2018. Advancing marine biological observations and data requirements of the complementary Essential Ocean Variables (EOVs) and Essential Biodiversity Variables (EBVs) frameworks. *Frontiers in Marine Science* 5:211, <https://doi.org/10.3389/fmars.2018.00211>.
- Naito, Y., D.P. Costa, T. Adachi, P.W. Robinson, S.H. Peterson, Y. Mitani, and A. Takahashi. 2017. Oxygen minimum zone: An important oceanographic habitat for deep-diving northern elephant seals, *Mirovanga angustirostris*. *Ecology and Evolution* 7:6,259–6,270, <https://doi.org/10.1002/ece3.3202>.
- NOAA (National Oceanographic and Atmospheric Administration). 2021a. California Current Integrated Ecosystem Assessment – CCIEA Publications: Reports, <https://www.integratedecosystemassessment.noaa.gov/regions/california-current/cc-publications-reports>.
- NOAA 2021b. California Current Integrated Ecosystem Assessment - Indicator Status Trends, <https://www.integratedecosystemassessment.noaa.gov/regions/california-current/cc-indicator-status-trends>.
- Newton, J.A., R.A. Feely, E.B. Jewett, P. Williamson, and J. Mathis. 2015. *Global Ocean Acidification Observing Network: Requirements and Governance Plan, 2nd ed.* GOA-ON, 61 pp., <https://archimer.ifremer.fr/doc/00651/76343/>.
- OBIS (Ocean Biodiversity Information System). 2021. <https://obis.org/>.
- Oestreich, W.K., J.A. Fahlbusch, D.E. Cade, J. Calambokidis, T. Margolina, J. Joseph, A.S. Friedlaender, M.F. McKenna, A.K. Stimpert, B.L. Southall, and others. 2020. Animal-borne metrics enable acoustic detection of blue whale migration. *Current Biology* 30(23):4,773–4,779, <https://doi.org/10.1016/j.cub.2020.08.105>.
- Ohman, M.D., R.E. Davis, J.T. Sherman, K.R. Grindley, B.M. Whitmore, C.F. Nickels, and J.S. Ellen. 2019. Zooglider: An autonomous vehicle for optical and acoustic sensing of zooplankton. *Limnology and Oceanography: Methods* 17 (1):69–86, <https://doi.org/10.1002/lom3.10301>.
- ONMS (Office of National Marine Sanctuaries). 2009. *Monterey Bay National Marine Sanctuary Condition Report 2009*. US Department of Commerce, National Oceanic and Atmospheric Administration, Office of National Marine Sanctuaries, Silver Spring, MD, 128 pp., <https://sanctuaries.noaa.gov/science/condition/mbnms/>.
- ONMS. 2015. *Monterey Bay National Marine Sanctuary Condition Report Partial Update: A New Assessment of the State of Sanctuary Resources 2015*. US Department of Commerce, National Oceanic and Atmospheric Administration, Office of National Marine Sanctuaries, Silver Spring, MD, 133 pp., <http://sanctuaries.noaa.gov/science/condition/monterey-bay-2015/>.
- ONMS. 2018. *Guide for Developing National Marine Sanctuary Condition Reports*. US Department of Commerce, National Oceanic and Atmospheric Administration, Office of National Marine Sanctuaries, Silver Spring, MD, 49 pp.
- Piatt, J.F., J.K. Parrish, H.M. Renner, S.K. Schoen, T.T. Jones, M.L. Arimitsu, K.J. Kuletz, B. Bodenstein, M. García-Reyes, R.S. Duerr, and others. 2020. Extreme mortality and reproductive failure of common murrelets resulting from the northeast

- Pacific marine heatwave of 2014–2016. *PLoS ONE* 15:e0226087, <https://doi.org/10.1371/journal.pone.0226087>.
- Ruhl, H.A., and K.L. Smith Jr. 2004. Shifts in deep-sea community structure linked to climate and food supply. *Science* 305:513–515, <https://doi.org/10.1126/science.1099759>.
- Ryan, J.P., R.M. Kudela, J.M. Birch, M. Blum, H.A. Bowers, F.P. Chavez, G.J. Doucette, K. Hayashi, R. Marin III, C.M. Mikulski, and others. 2017. Causality of an extreme harmful algal bloom in Monterey Bay, California, during the 2014–2016 northeast Pacific warm anomaly. *Geophysical Research Letters* 44(11):5711–5719, <https://doi.org/10.1002/2017GL072637>.
- Ryan, J.P., D.E. Cline, J.E. Joseph, T. Margolina, J.A. Santora, R.A. Kudela, F.P. Chavez, J.T. Pennington, C. Wahl, R. Michisaki, and others. 2019. Humpback whale song occurrence reflects ecosystem variability in feeding and migratory habitat of the northeast Pacific. *PLoS ONE* 14(9):e0222456, <https://doi.org/10.1371/journal.pone.0222456>.
- Ryan, J.P., J.E. Joseph, T. Margolina, L.T. Hatch, A. Azzara, A. Reyes, B.L. Southall, A. DeVogelaere, L.E. Peavey, D.E. Cline, and others. 2021. Reduction of low-frequency vessel noise in Monterey Bay National Marine Sanctuary during the COVID-19 pandemic. *Frontiers in Marine Science* 8:656566, <https://doi.org/10.3389/fmars.2021.656566>.
- Sakuma, K.M., J.C. Field, N.J. Mantua, S. Ralston, B.B. Marinovic, and C.N. Carrion. 2016. Anomalous epipelagic micronekton assemblage patterns in the neritic waters of the California current in spring 2015 during a period of extreme ocean conditions. *CalCOFI Reports* 57:163–183, http://calcofi.org/publications/calcofireports/v57/Vol57-Sakuma_pages.163-183.pdf.
- Samhuri, J.F., A.J. Haupt, P.S. Levin, J.S. Link, and R. Shuford. 2014. Lessons learned from developing integrated ecosystem assessments to inform marine ecosystem-based management in the USA. *ICES Journal of Marine Science* 71(5):1,205–1,215, <https://doi.org/10.1093/icesjms/fst141>.
- Samhuri, J.F., K.S. Andrews, G. Fay, C.J. Harvey, E.L. Hazen, S.M. Hennessey, K. Holsman, M.E. Hunsicker, S.I. Large, K.N. Marshall, and others. 2017. Defining ecosystem thresholds for human activities and environmental pressures in the California Current. *Ecosphere* 8(6):e01860, <https://doi.org/10.1002/ecs2.1860>.
- Santora, J.A., E.L. Hazen, I.D. Schroeder, S.J. Bograd, K.M. Sakuma, and J.C. Field. 2017. Impacts of ocean-climate variability on biodiversity of pelagic forage species in an upwelling ecosystem. *Marine Ecology Progress Series* 580:205–220, <https://doi.org/10.3354/meps12278>.
- Santora, J.A., N.J. Mantua, I.D. Schroeder, J.C. Field, E.L. Hazen, S.J. Bograd, W.J. Sydeman, B.K. Well, J. Calambokidis, L. Saez, and others. 2020. Habitat compression and ecosystem shifts as potential links between marine heatwave and record whale entanglements. *Nature Communications* 11:536, <https://doi.org/10.1038/s41467-019-14215-w>.
- Santora, J.A., I.D. Schroeder, S.J. Bograd, F.P. Chavez, M.A. Cimino, J. Fiechter, E.L. Hazen, M.T. Kavanaugh, M. Messié, R.R. Miller, and others. 2021. Pelagic biodiversity, ecosystem function, and services: An integrated observing and modeling approach. *Oceanography* 34(2):16–37, <https://doi.org/10.5670/oceanog.2021.212>.
- Schwing, F.B., T. Murphree, and P.M. Green. 2002. The Northern Oscillation Index (NOI): A new climate index for the northeast Pacific. *Progress in Oceanography* 53:115–139, [https://doi.org/10.1016/S0079-6611\(02\)00027-7](https://doi.org/10.1016/S0079-6611(02)00027-7).
- Sequeira, A.M.M., J.P. Rodríguez, V.M. Eguíluz, R. Harcourt, M. Hindell, D.W. Sims, C.M. Duarte, D.P. Costa, J. Fernández-Gracia, L.C. Ferreira, and others. 2018. Convergence of marine megafauna movement patterns in coastal and open oceans. *Proceedings of the National Academy Sciences of the United States of America* 115:3,072–3,077, <https://doi.org/10.1073/pnas.1716137115>.
- Simonis, A.E., K.A. Forney, S. Rankin, J. Ryan, Y. Zhang, A. DeVogelaere, J. Joseph, T. Margolina, A. Krumpel, and S. Baumann-Pickering. 2020. Seal bomb noise as a potential threat to Monterey Bay harbor porpoise. *Frontiers in Marine Science* 7:142, <https://doi.org/10.3389/fmars.2020.00142>.
- Simons, R.A. 2019. ERDDAP, <https://coastwatch.pfeg.noaa.gov/erddap>.
- Sosik, H.M., and R.J. Olson. 2007. Automated taxonomic classification of phytoplankton sampled with imaging-in-flow cytometry. *Limnology and Oceanography: Methods* 5:204–216, <https://doi.org/10.4319/lom.2007.5.204>.
- Southall, B., J. Berkson, D. Bowen, R. Brake, J. Eckman, J. Field, R. Gisiner, S. Gregerson, W. Lang, J. Lewandoski, and others. 2009. *Addressing the Effects of Human-Generated Sound on Marine Life: An Integrated Research Plan for US Federal Agencies*. Interagency Task Force on Anthropogenic Sound and the Marine Environment of the Joint Subcommittee on Ocean Science and Technology, National Oceanic and Atmospheric Administration, Washington, DC, 72 pp., <https://tethys.pnnl.gov/publications/addressing-effects-human-generated-sound-marine-life-integrated-research-plan-us>.
- Spector, P., B.D. Best, J. Ranganathan, T. Murray, J.A. Brown, C. Caldwell, G. Canonico, and A. DeVogelaere. 2021. *Webenizing Condition Reports: Communicating Data-Driven Ecosystem Indicators in a Visually Engaging and Interactive Online Platform*. National Marine Sanctuaries Conservation Series ONMS-21-11. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Office of National Marine Sanctuaries.
- TDWG (Biodiversity Information Standards). 2021. Darwin Core, <https://www.tdwg.org/standards/dwcl/>.
- Wieczorek, J., D. Bloom, R. Guralnick, S. Blum, M. Döring, R. Giovanni, T. Robertson, and D. Vieglais. 2012. Darwin Core: An evolving community-developed biodiversity data standard. *PLoS ONE* 7(1):e29715, <https://doi.org/10.1371/journal.pone.0029715>.
- Wilkinson, M.D., M. Dumontier, I.J.J. Aalbersberg, G. Appleton, M. Axton, A. Baak, N. Blomberg, J.-W. Boiten, L.B. da Silva Santos, P.E. Bourne, and others. 2016. The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* 3:160018, <https://doi.org/10.1038/sdata.2016.18>.
- Woodson, C.B., M.A. McManus, J.A. Tyburczy, J.A. Barth, L. Washburn, J.E. Caselle, M.H. Carr, D.P. Malone, P.T. Raimondi, B.A. Menge, and S.R. Palumbi. 2012. Coastal fronts set recruitment and connectivity patterns across multiple taxa. *Limnology and Oceanography* 57:582–596, <https://doi.org/10.4319/lo.2012.57.2.0582>.
- WoRMS Editorial Board. 2021. World Register of Marine Species, <http://www.marinespecies.org>.
- Grant# 80NSSC20M0001), and “Implementing a Marine Biodiversity Observation Network (MBON) in South Florida to Advance Ecosystem-Based Management” (NASA Grant # 80NSSC20K0017; ONR/NOAA/US IOOS grant NA19NOS0120199), awarded through the National Ocean Partnership Program (NOPP). NOPP includes contributions from BOEM, NASA, ONR, and NOAA. Thanks to Steve Gittings and Gabrielle Canonico for reviewing drafts of this manuscript.

AUTHORS

Henry A. Ruhl (hruhl@mbari.org) is CeNCOOS Director, Monterey Bay Aquarium Research Institute (MBARI), Moss Landing, CA, USA. **Jennifer A. Brown** is Principal, ECOS Consulting LLC, Lafayette, CA, USA, and Ecosystem Scientist, National Oceanic and Atmospheric Administration (NOAA), Monterey Bay National Marine Sanctuary (MBNMS), Monterey, CA, USA. **Alexandra R. Harper** is CeNCOOS Program Manager, MBARI, Moss Landing, CA, USA. **Elliott L. Hazen** is Research Ecologist, NOAA, Southwest Fisheries Science Center (SWFSC), Monterey, CA, USA. **Lynn deWitt** is IT Specialist, NOAA, SWFSC, Santa Cruz, CA, USA. **Patrick Daniel** is CeNCOOS Data Manager, MBARI, Moss Landing, CA, USA. **Andrew DeVogelaere** is Research Coordinator, NOAA, MBNMS, Monterey, CA, USA. **Raphael M. Kudela** is Professor, University of California Santa Cruz (UCSC), Santa Cruz, CA, USA. **John P. Ryan** is Senior Research Specialist, MBARI, Moss Landing, CA, USA. **Alexis D. Fischer** was Postdoctoral Fellow, UCSC, Santa Cruz, CA, USA, and is currently Associate Scientist, University Corporation for Atmospheric Research, and Visiting Scientist, NOAA, Northwest Fisheries Science Center, Seattle, WA, USA. **Frank E. Muller-Karger** is Professor and Director, IMaRS, College of Marine Science, University of South Florida, St. Petersburg, FL, USA. **Francisco P. Chavez** is Senior Scientist, MBARI, Moss Landing, CA, USA.

ARTICLE CITATION

Ruhl, H.A., J.A. Brown, A.R. Harper, E.L. Hazen, L. deWitt, P. Daniel, A. DeVogelaere, R.M. Kudela, J.P. Ryan, A.D. Fischer, F.E. Muller-Karger, and F.P. Chavez. 2021. Integrating biodiversity and environmental observations in support of national marine sanctuary and large marine ecosystem assessments. *Oceanography* 34(2):142–155, <https://doi.org/10.5670/oceanog.2021.221>.

COPYRIGHT & USAGE

This is an open access article made available under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution, and reproduction in any medium or format as long as users cite the materials appropriately, provide a link to the Creative Commons license, and indicate the changes that were made to the original content.

ACKNOWLEDGMENTS

This work was made possible in part by efforts of NOAA, including the Integrated Ecosystem Assessment, the Office of National Marine Sanctuaries, the National Marine Fisheries Service, Integrated Ocean Observing System, and Marine Biodiversity Observation Network. Support funding included the project “National Marine Sanctuaries as Sentinel Sites for a Demonstration Marine Biodiversity Observation Network (Sanctuaries MBON)” (NASA Grant #NNX14AP62A) and its follow-up projects “The CeNCOOS MBON: Integrating remote sensing, in situ data and models to understand central California ecosystem responses to environmental change” (NASA