

SIDEBAR > OBSERVATIONS OF DECLINING PRIMARY PRODUCTIVITY IN THE WESTERN BERING STRAIT

By Karen E. Frey, Jaclyn Clement Kinney, Larry V. Stock, and Robert Osinski

The shallow (~50 m deep), narrow (~85 km wide) Bering Strait is the sole marine link between the Pacific and Arctic Oceans and represents a critical northward throughflow of freshwater, nutrients, and heat into Arctic waters from lower latitudes (Woodgate and Peralta-Ferriz, 2021). Three water masses enter the Chukchi Sea through the Bering Strait from the Pacific: Anadyr Water (AW), Bering Shelf Water (BSW), and Alaskan Coastal Water (ACW) (Coachman et al., 1975). The western Bering Strait in particular has long been known to be a region of consistently high primary productivity throughout the spring and summer open-water season (Sambrotto et al., 1984; Springer and McRoy, 1993; Brown et al., 2011). This productivity is sustained through the delivery of high-nutrient AW waters via the northern branch of the bifurcated Bering Slope Current (Clement Kinney et al., 2009, 2022; Lowry et al., 2015; Pickart et al., 2016) that also causes the Chukchi Sea to the north to be one of the most productive shelves in the Arctic (Hill et al., 2018). Western Bering Strait waters are clearly differentiated from lower productivity waters observed in the eastern Bering Strait that are characterized by relatively low-nutrient, freshwater-dominated ACW (Woodgate and Aagaard, 2005; Lee et al., 2007). However, time series of satellite observations over the last two decades have revealed statistically significant early season (June) declining trends in chlorophyll-*a* concentrations and primary productivity in the western Bering Strait. In particular, June chlorophyll-*a* concentrations have declined by ~58%, and June primary productivity has declined by ~34% over the 2003–2020 period. These declining trends appear to be associated with reductions in sea ice cover and increases in primary production upstream in the Gulf of Anadyr during May, with potential implications for decreased nutrient availability downstream in the western Bering Strait during June.

To investigate recent biological change in the Bering Strait, we compiled a satellite-based time series of chlorophyll-*a* concentrations derived from Aqua-Moderate Resolution Imaging Spectroradiometer (Aqua-MODIS) calibrated radiances using two algorithms: the OC3m algorithm that was developed at NASA Goddard Space Flight Center (GSFC) and makes use of band ratios and in situ measurements (O'Reilly et al., 1998) and the CI algorithm that makes use of reflectance differences in conjunction with a model (Hu et al., 2012). The data are made available by the Ocean Biology Processing Group and were downloaded from the GSFC Distributed Active Archive Center (DAAC) at <https://oceandata.sci.gsfc.nasa.gov/MODIS-Aqua/Mapped/>

[Daily/4km/chlor_a/](#). Chlorophyll-*a* concentration data were also combined with sea surface temperature data and additional data sets to derive net primary productivity using a broadly utilized algorithm (Behrenfeld and Falkowski, 1997) that has previously been employed to report changes across the Arctic region (Frey et al., 2021). Monthly chlorophyll-*a* and primary productivity data were only utilized where sea ice concentrations were <10% and were otherwise reported as missing data. For further context, we investigated sea ice concentration data obtained from the Special Sensor Microwave/Imager (SSM/I) and Special Sensor Microwave Imager/Sounder (SSMIS) passive microwave instruments, calculated using the Goddard Bootstrap (SB2) algorithm (Comiso et al., 2017a,b). Modeled surface nitrate concentrations were obtained from the Regional Arctic System Model (RASM; e.g., Clement Kinney et al., 2020). For all data sets (chlorophyll-*a*, primary productivity, sea ice, and surface nitrate), monthly time series were compiled for May and June, and the Theil-Sen median decadal trends for each month (2003–2020) were calculated, with statistically significant ($p < 0.1$) trends identified using the non-parametric Mann-Kendall test for monotonic trend (Mann, 1945; Kendall, 1975). The Theil-Sen median trend uses a robust non-parametric trend operator that is particularly well suited for assessing the rate of change in noisy and/or short time series (Hoaglin et al., 2000), which in this study is 18 years. For those data sets that include missing data (chlorophyll-*a* and primary productivity), we show only those trends for pixels that had at least 71% of the time series present (or in the case of this study, 13 of the 18 time steps). This requirement ensures that only robust trends are reported, given that the “breakdown bound” for the Theil-Sen trend is 29% (meaning that unknown or potentially “wild” values would have to persist for more than 29% of a time series in order to affect the overall trend values; Hoaglin et al., 2000).

Increasing trends in marine primary productivity across the Arctic owing to shifts in sea ice cover, seawater temperatures, and nutrient availability have been widely reported (Arrigo et al., 2008; Pabi et al., 2008; Arrigo and van Dijken, 2015; Clement Kinney et al., 2020; Lewis et al., 2020; Frey et al., 2021). In contrast to those reports of large-scale increases in primary productivity, [Figure 1](#) identifies an important and unusual regional location of early season (June) declines in productivity, with potential implications for nutrient and carbon delivery downstream (northward) across the Chukchi Sea shelf. During May (over the 2003–2020 period), we observe strong declines in sea ice concentration in the Gulf of Anadyr

(Figure 1a) with increases in surface nitrate concentrations (Figure 1c), and these are in turn associated with increasing trends in both chlorophyll-*a* (Figure 1e) and primary productivity (Figure 1g). However, during June, we observe (and model with RASM, not shown) strikingly strong and spatially cohesive declines in chlorophyll-*a* (Figure 1f) and primary productivity (Figure 1h) downstream of the Gulf of Anadyr in the western Bering Strait along the coast of the Chukotka Peninsula. By June, sea ice has typically already exhibited seasonal breakup in the Bering Strait region (Frey et al., 2015), and we see no trends in sea ice cover in the western Bering Strait (Figure 1b). Significant declining trends in June surface nitrate concentrations (Figure 1d) geographically mirror the observed declines in chlorophyll-*a* (Figure 1f) and primary productivity (Figure 1h). It is important to note that the potential for increased presence of subsurface chlorophyll maxima (as a result of deepening nutriclines) may be challenging to quantify seasonally via satellite data in the Chukchi Sea (Arrigo et al., 2011; Ardyna et al., 2013; Brown et al., 2015). Nonetheless, we hypothesize that because of the May declines of sea ice in the Gulf of Anadyr and resulting increases in May chlorophyll-*a*/primary production in that region, available nutrients downstream in the western Bering Strait during June are depleted, and chlorophyll-*a*/primary productivity therefore have declined over time there as well. In particular, in the western Bering Strait (within the region designated as statistically significant for June chlorophyll-*a* concentrations; Figure 1f), June chlorophyll-*a* concentrations have changed by approximately -58% (from 4.2 mg/m^3 to 1.8 mg/m^3), and June primary productivity has changed by approximately -34% (from $2,418 \text{ mg C/m}^2/\text{day}$ to $1,606 \text{ mg C/m}^2/\text{day}$). These shifts represent chlorophyll-*a* trends of $-1.52 \text{ mg/m}^3/\text{decade}$ and primary productivity trends of $-477.8 \text{ mg C/m}^2/\text{day}/\text{decade}$. However, increases in chlorophyll-*a* and primary productivity in the western Bering Strait primarily during September (not shown) counteract these June decreases, so overall annual primary productivity rates in this region are not significant. Thus, while annual productivity may not have changed substantially, observed shifts in the seasonal distribution of productivity may indeed have




FIGURE 1. Decadal Theil-Sen median trends for May/June over the years 2003–2020 in (a,b) sea ice concentrations, (c,d) surface nitrate concentrations, (e,f) chlorophyll-*a* concentrations, and (g,h) primary productivity. Hatched regions indicate statistically significant ($p < 0.1$) trends, determined using the Mann-Kendall test for trend. Distributed Biological Observatory (DBO) sites 1, 2, and 3 (Grebmeier et al., 2019) are shown for geographic context.

profound consequences for marine ecosystem functioning across this region.

Despite measurements of overall, large-scale increases in primary productivity across the Arctic Ocean over recent decades, heterogeneity in shifts of nutrient availability to upper ocean waters across the region has also led to a spatial mosaic of both increases and decreases in productivity (Juraneck, 2022, in this issue). For example, while earlier sea ice retreat can result in stronger blooms in Arctic shelf regions, increased sea ice melt can also result in reduced production in portions of the central Arctic owing to enhanced stratification (Song et al., 2021). Furthermore, moored sensor-based measurements of dissolved inorganic nitrogen (DIN) in bottom waters in the northern Bering Sea indicate high inter-annual variability but an overall decline of ~50% over the 2005–2017 period, with strong correlations of late summer/early fall DIN resulting in primary productivity downstream on the northern Chukchi shelf the following May (Mordy et al., 2020). Likewise, the early season declines in primary productivity in the western Bering Strait found in this study should undoubtedly have important consequences for the further downstream delivery of carbon and otherwise excess nutrients to the Herald Canyon and western/central Chukchi Shelf regions, important hotspots for biological productivity in the Arctic (Arrigo et al., 2012, 2014; Linders et al., 2017; Li et al., 2019). Changes in the seasonal and spatial distribution of spring phytoplankton blooms in the Pacific Arctic will also likely have important effects on pelagic-benthic coupling in a region with historically high benthic biomass and large populations of seabirds and marine mammals that depend upon benthic prey for survival (Grebmeier et al., 2006, 2018).

The observations of change in the western Bering Strait reported here provide an important example of the heterogeneity of ecosystem responses to climate change, where primary productivity does not always increase with declines in sea ice cover. Moreover, it is important to consider how environmental changes such as sea ice decline can have vital impacts on ecosystem functioning not only locally but also through resulting impacts on nutrient delivery downstream along a conveyor belt system of ocean currents. 

REFERENCES

- Ardyna, M., M. Babin, M. Gosselin, E. Devred, S. Bélanger, A. Matsuoka, and J.-É. Tremblay. 2013. Parameterization of vertical chlorophyll σ in the Arctic Ocean: Impact of the subsurface chlorophyll maximum on regional, seasonal, and annual primary production estimates. *Biogeosciences* 10(6):4,383–4,404, <https://doi.org/10.5194/bg-10-4383-2013>.
- Arrigo, K.R., G. van Dijken, and S. Pabi. 2008. Impact of a shrinking Arctic ice cover on marine primary production. *Geophysical Research Letters* 35(19), <https://doi.org/10.1029/2008GL035028>.
- Arrigo, K.R., P.A. Matrai, and G.L. van Dijken. 2011. Primary productivity in the Arctic Ocean: Impacts of complex optical properties and subsurface chlorophyll maxima on large-scale estimates. *Journal of Geophysical Research: Oceans* 116(C11), <https://doi.org/10.1029/2011JC007273>.
- Arrigo, K.R., D.K. Perovich, R.S. Pickart, Z.W. Brown, G.L. van Dijken, K.E. Lowry, M.M. Mills, M.A. Palmer, W.M. Balch, and F. Bahr. 2012. Massive phytoplankton blooms under Arctic sea ice. *Science* 336(6087):1,408–1,408, <https://doi.org/10.1126/science.1215065>.
- Arrigo, K.R., D.K. Perovich, R.S. Pickart, Z.W. Brown, G.L. van Dijken, K.E. Lowry, M.M. Mills, M.A. Palmer, W.M. Balch, and N.R. Bates. 2014. Phytoplankton blooms beneath the sea ice in the Chukchi Sea. *Deep Sea Research Part II* 105:1–16, <https://doi.org/10.1016/j.dsr2.2014.03.018>.
- Arrigo, K.R., and G.L. van Dijken. 2015. Continued increases in Arctic Ocean primary production. *Progress in Oceanography* 136:60–70, <https://doi.org/10.1016/j.pocean.2015.05.002>.
- Behrenfeld, M.J., and P.G. Falkowski. 1997. Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnology and Oceanography* 42(1):1–20, <https://doi.org/10.4319/lo.1997.42.1.0001>.
- Brown, Z.W., G.L. van Dijken, and K.R. Arrigo. 2011. A reassessment of primary production and environmental change in the Bering Sea. *Journal of Geophysical Research: Oceans* 116(C8), <https://doi.org/10.1029/2010JC006766>.
- Brown, Z.W., K.E. Lowry, M.A. Palmer, G.L. van Dijken, M.M. Mills, R.S. Pickart, and K.R. Arrigo. 2015. Characterizing the subsurface chlorophyll a maximum in the Chukchi Sea and Canada Basin. *Deep Sea Research Part II* 118:88–104, <https://doi.org/10.1016/j.dsr2.2015.02.010>.
- Clement Kinney, J., W. Maslowski, and S. Okkonen. 2009. On the processes controlling shelf-basin exchange and outer shelf dynamics in the Bering Sea. *Deep Sea Research Part II* 56(17):1,351–1,362, <https://doi.org/10.1016/j.dsr2.2008.10.023>.
- Clement Kinney, J., W. Maslowski, R. Osinski, M. Jin, M. Frants, N. Jeffery, and Y.J. Lee. 2020. Hidden production: On the importance of pelagic phytoplankton blooms beneath Arctic sea ice. *Journal of Geophysical Research: Oceans* 125(9):e2020JC016211, <https://doi.org/10.1029/2020JC016211>.
- Clement Kinney, J., K.M. Assmann, W. Maslowski, G. Björk, M. Jakobsson, S. Jutterström, Y.J. Lee, R. Osinski, I. Semiletov, and A. Ulfso. 2022. On the circulation, water mass distribution, and nutrient concentrations of the western Chukchi Sea. *Ocean Science* 18:29–49, <https://doi.org/10.5194/os-18-29-2022>.
- Coachman, L.K., K. Aagaard, and R. Tripp. 1975. *Bering Strait: The Regional Physical Oceanography*. University of Washington Press.
- Comiso, J.C., R.A. Gersten, L.V. Stock, J. Turner, G.J. Perez, and K. Cho. 2017a. Positive trend in the Antarctic sea ice cover and associated changes in surface temperature. *Journal of Climate* 30(6):2,251–2,267, <https://doi.org/10.1175/JCLI-D-16-0408.1>.
- Comiso, J.C., W.N. Meier, and R. Gersten. 2017b. Variability and trends in the Arctic Sea ice cover: Results from different techniques. *Journal of Geophysical Research: Oceans* 122(8):6,883–6,900, <https://doi.org/10.1002/2017JC012768>.
- Frey, K.E., G. Moore, L.W. Cooper, and J.M. Grebmeier. 2015. Divergent patterns of recent sea ice cover across the Bering, Chukchi, and Beaufort seas of the Pacific Arctic Region. *Progress in Oceanography* 136:32–49, <https://doi.org/10.1016/j.pocean.2015.05.009>.
- Frey, K.E., J. Comiso, L. Cooper, J. Grebmeier, and L. Stock. 2021. Arctic ocean primary productivity: The response of marine algae to climate warming and sea ice decline. Pp. 46–57 in *Arctic Report Card 2021*. T.A. Moon, M.L. Druckenmiller, and R.L. Thoman, eds, <https://doi.org/10.25923/kxhb-dw16>.
- Grebmeier, J.M., J.E. Overland, S.E. Moore, E.V. Farley, E.C. Carmack, L.W. Cooper, K.E. Frey, J.H. Helle, F.A. McLaughlin, and S.L. McNutt. 2006. A major ecosystem shift in the northern Bering Sea. *Science* 311(5766):1,461–1,464, <https://doi.org/10.1126/science.1121365>.
- Grebmeier, J.M., K.E. Frey, L.W. Cooper, and M. Kędra. 2018. Trends in benthic macrofaunal populations, seasonal sea ice persistence, and bottom water temperatures in the Bering Strait region. *Oceanography* 31(2):136–151, <https://doi.org/10.5670/oceanog.2018.224>.
- Grebmeier, J.M., S.E. Moore, L.W. Cooper, and K.E. Frey. 2019. The Distributed Biological Observatory: A change detection array in the Pacific Arctic—An introduction. *Deep Sea Research Part II* 162:1–7, <https://doi.org/10.1016/j.dsr2.2019.05.005>.
- Hill, V., M. Ardyna, S.H. Lee, and D.E. Varela. 2018. Decadal trends in phytoplankton production in the Pacific Arctic Region from 1950 to 2012. *Deep Sea Research Part II* 152:82s94, <https://doi.org/10.1016/j.dsr2.2016.12.015>.
- Hoaglin, D., F. Mosteller, and J. Tukey, eds. 2000. *Understanding Robust and Exploratory Data Analysis*. Wiley Classics Library, Wiley, New York, 472 pp.
- Hu, C., Z. Lee, and B. Franz. 2012. Chlorophyll a algorithms for oligotrophic oceans: A novel approach based on three-band reflectance difference. *Journal of Geophysical Research: Oceans* 117(C1), <https://doi.org/10.1029/2011JC007395>.

- Juranek, L.W. 2022. Changing biogeochemistry of the Arctic Ocean: Surface nutrient and CO₂ cycling in a warming, melting north. *Oceanography* 35(3–4):144–155, <https://doi.org/10.5670/oceanog.2022.120>.
- Kendall, M. 1975. *Rank Correlation Methods*, 4th ed. Griffen, London 202 pp.
- Lee, S.H., T.E. Whitledge, and S.-H. Kang. 2007. Recent carbon and nitrogen uptake rates of phytoplankton in Bering Strait and the Chukchi Sea. *Continental Shelf Research* 27(17):2,231–2,249, <https://doi.org/10.1016/j.csr.2007.05.009>.
- Lewis, K., G. van Dijken, and K.R. Arrigo. 2020. Changes in phytoplankton concentration now drive increased Arctic Ocean primary production. *Science* 369(6500):198–202, <https://doi.org/10.1126/science.aay8380>.
- Li, M., R.S. Pickart, M.A. Spall, T.J. Weingartner, P. Lin, G. Moore, and Y. Qi. 2019. Circulation of the Chukchi Sea shelfbreak and slope from moored timeseries. *Progress in Oceanography* 172:14–33, <https://doi.org/10.1016/j.pocean.2019.01.002>.
- Linders, J., R.S. Pickart, G. Björk, and G. Moore. 2017. On the nature and origin of water masses in Herald Canyon, Chukchi Sea: Synoptic surveys in summer 2004, 2008, and 2009. *Progress in Oceanography* 159:99–114, <https://doi.org/10.1016/j.pocean.2017.09.005>.
- Lowry, K.E., R.S. Pickart, M.M. Mills, Z.W. Brown, G.L. van Dijken, N.R. Bates, and K.R. Arrigo. 2015. The influence of winter water on phytoplankton blooms in the Chukchi Sea. *Deep Sea Research Part II* 118:53–72, <https://doi.org/10.1016/j.dsr2.2015.06.006>.
- Mann, H. 1945. Non-parametric test against trend. *Econometrica* 13:245–259, <https://doi.org/10.2307/1907187>.
- Mordy, C.W., S. Bell, E.D. Cokolet, C. Ladd, G. Lebon, P. Proctor, P. Stabeno, D. Strausz, E. Wisegarver, and K. Wood. 2020. Seasonal and interannual variability of nitrate in the eastern Chukchi Sea: Transport and winter replenishment. *Deep Sea Research Part II* 177:104807, <https://doi.org/10.1016/j.dsr2.2020.104807>.
- O'Reilly, J.E., S. Maritorena, B.G. Mitchell, D.A. Siegel, K.L. Carder, S.A. Garver, M. Kahru, and C. McClain. 1998. Ocean color chlorophyll algorithms for SeaWiFS. *Journal of Geophysical Research: Oceans* 103(C11):24,937–24,953, <https://doi.org/10.1029/98JC02160>.
- Pabi, S., G.L. van Dijken, and K.R. Arrigo. 2008. Primary production in the Arctic Ocean, 1998–2006. *Journal of Geophysical Research: Oceans* 113(C8), <https://doi.org/10.1029/2007JC004578>.
- Pickart, R.S., G. Moore, C. Mao, F. Bahr, C. Nobre, and T.J. Weingartner. 2016. Circulation of winter water on the Chukchi shelf in early summer. *Deep Sea Research Part II* 130:56–75, <https://doi.org/10.1016/j.dsr2.2016.05.001>.
- Sambrotto, R., J. Goering, and C. McRoy. 1984. Large yearly production of phytoplankton in the western Bering Strait. *Science* 225(4667):1,147–1,150, <https://doi.org/10.1126/science.225.4667.1147>.
- Song, H., R. Ji, M. Jin, Y. Li, Z. Feng, Ø. Varpe, and C.S. Davis. 2021. Strong and regionally distinct links between ice-retreat timing and phytoplankton production in the Arctic Ocean. *Limnology and Oceanography* 66(6):2,498–2,508, <https://doi.org/10.1002/lno.11768>.
- Springer, A.M., and C.P. McRoy. 1993. The paradox of pelagic food webs in the northern Bering Sea: Part III. Patterns of primary production. *Continental Shelf Research* 13(5–6):575–599, [https://doi.org/10.1016/0278-4343\(93\)90095-F](https://doi.org/10.1016/0278-4343(93)90095-F).
- Woodgate, R.A., and K. Aagaard. 2005. Revising the Bering Strait freshwater flux into the Arctic Ocean. *Geophysical Research Letters* 32(2), <https://doi.org/10.1029/2004GL021747>.
- Woodgate, R.A., and C. Peralta-Ferriz. 2021. Warming and freshening of the Pacific inflow to the Arctic from 1990–2019 implying dramatic shoaling in Pacific Winter Water ventilation of the Arctic water column. *Geophysical Research Letters* 48(9):e2021GL092528, <https://doi.org/10.1029/2021GL092528>.

ACKNOWLEDGMENTS

K. Frey acknowledges financial support from the US National Science Foundation (NSF) Arctic Observing Network (AON) Program (grant number 1917434). J. Clement Kinney acknowledges financial support from the US Department of Energy (Office of Science, Office of Basic Energy Sciences and Energy Efficiency and Renewable Energy, Solar Energy Technology Program; grant number RGMA IAA#DE-SC0014117) and NSF (grant number GEO/PLR ARCSS IAA#1417888). L. Stock is grateful for the support provided by the NASA Ocean Biology and Biogeochemistry Program. R. Osinski was supported by the Ministry of Science and Higher Education in Poland under international project agreement number 3808/FAO/2017/O RASMer.

AUTHORS

Karen E. Frey (kfrey@clarku.edu) is Professor, Graduate School of Geography, Clark University, Worcester, MA, USA. **Jaclyn Clement Kinney** is Research Associate Professor, Naval Postgraduate School, Monterey, CA, USA. **Larry V. Stock** is Scientific Programmer, Cryospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA. **Robert Osinski** is a researcher at the Institute of Oceanology, Polish Academy of Sciences, Sopot, Poland.

ARTICLE CITATION

Frey, K.E., J. Clement Kinney, L.V. Stock, and R. Osinski. 2022. Observations of declining primary productivity in the western Bering Strait. *Oceanography* 35(3–4):140–143, <https://doi.org/10.5670/oceanog.2022.123>.